

•

Burning Trends and Potential Negative Effects of Suppressing Wetland Fires on Flatwoods Salamanders

David C. Bishop¹
Carola A. Haas

Department of Fisheries and
Wildlife Sciences
Virginia Polytechnic Institute
and State University
Blacksburg, VA 24061, USA

•

¹ Corresponding author:
dabishop@vt.edu

ABSTRACT: The federally threatened flatwoods salamander (*Ambystoma cingulatum* Cope) inhabits fire-adapted longleaf pine (*Pinus palustris* Miller) flatwoods and savannas in the southeastern coastal plain. It breeds in ephemeral wetlands that typically dry during summer and refill in fall and winter. We collected data on burning trends of *A. cingulatum* habitat across the range of the species. Rangewide surveys indicate that prescribed fires typically are applied in winter and early spring. If prescribed fires are conducted primarily in winter, fires are unlikely to burn through water-filled wetlands, potentially affecting larval habitat. Based on what is known about the natural history of the species, the historical burning regime of the longleaf pine ecosystem, and the effects of fires on ephemeral wetlands, we suggest that land managers diversify their fire-management strategy to increase the likelihood of burning the breeding wetlands of flatwoods salamanders.

Index terms: *Ambystoma cingulatum*, amphibian, fire, longleaf, wetland

INTRODUCTION

The impacts of forest management activities on amphibian communities have received more attention in recent years. Most studies have focused on the negative effects of timber harvesting on amphibian populations (e.g., Petranka et al. 1994, Ash 1997, Grialou et al. 2000, Knapp et al. 2003). But other activities, such as ditching and bedding (often used in coastal plain silvicultural operations (Pechmann et al. 1989, Means et al. 1996)) and fire suppression, may also affect amphibian diversity and abundance (deMaynadier and Hunter 1995, Russell et al. 1999, Pilliod et al. 2003). The federally threatened flatwoods salamander (Ambystomatidae: *Ambystoma cingulatum* Cope) has declined throughout its range. Major threats to the species include the degradation and loss of habitat from development, agriculture and silviculture, modification of breeding wetland hydrology, and fire suppression (Means et al. 1996, Palis 1996, U.S. Fish and Wildlife Service 1999).

Flatwoods salamanders inhabit fire-adapted longleaf pine (*Pinus palustris* Miller) flatwoods and savannas in the southeastern coastal plain. They breed in ephemeral wetlands (Means 1972, Anderson and Williamson 1976, Palis 1997a), which typically dry in summer and refill in fall and winter (Palis 1997b, Stevenson, unpubl. data). The historical fire regime of the longleaf pine included frequent summer fires from both lightning strikes and Native Americans (Ware et al. 1993, Battle and Golladay 2003). Winter fires are less likely to move through wetland basins than summer fires, except in drought years. Fire exclusion may

impact vegetation composition, solar radiation, water temperature, water chemistry, hydrology, nutrient cycling, and productivity of wetlands (Kirkman 1995, Battle and Golladay 2003, Pilliod et al. 2003).

The purpose of this paper is to relate what is known about the natural history of flatwoods salamanders to the potential role of fire in structuring their breeding wetlands. We surveyed public lands containing known breeding populations of *A. cingulatum* and summarized burning patterns across the range of the species. Because data are limited on the effects of burning regime on flatwoods salamanders, we suggest fire-management decisions be based on what is known about the natural history of the species, the historical burning regime of the longleaf pine ecosystem, and the effects of fires on ephemeral wetlands.

METHODS

We collected fire histories from state and federal lands containing *A. cingulatum* breeding populations across the range of the species. Participants (Table 1) provided months and years of fires (Total N = 228) for an 11-year period (1 January 1992 to 31 December 2002). These data were used to evaluate the frequency of fires in surrounding uplands, not whether fires actually burned through wetlands, as these data were not available. However, because wetlands typically are burned in conjunction with surrounding uplands, these data can be used to assess the likelihood that fires would burn through wetland basins. We did not have sufficient data to separate wildfires from prescribed fires.

Table 1. Locations that contributed fire histories. The number of breeding wetlands (N = 154) indicates the number of *Ambystoma cingulatum* breeding wetlands for which we were given data.

Location	State	Breeding Wetlands	Fire History
Apalachicola National Forest	FL	50	1992–2002
Eglin Air Force Base	FL	18	1992–2002
Fort Stewart (Army)	GA	18	1992–2002
Francis Marion National Forest	SC	5	1992–2002
Holley Field (Navy)	FL	3	1992–2002
Hurlburt Field (Air Force)	FL	13	1992–2002
J. W. Jones Research Center (Univ. Georgia)	GA	4	1993–2002
Mayhaw Wildlife Management Area	GA	1	1992–2002
Pine Log State Forest	FL	1	1992–2002
St. Mark's National Wildlife Refuge	FL	41	1992–2002

fronts and decreasing atmospheric pressure (Means 1972, Anderson and Williamson 1976, Palis 1997a, Safer 2001). They typically breed in open-canopy wetlands (Anderson and Williamson 1976, Palis 1997b). Larvae can be captured December–April (Means 1972, Palis 1996) by dipnetting herbaceous vegetation (Palis 1996; Sekerak et al. 1996, pers. observations), which typically is more abundant in open-canopy wetlands and in open sections within a wetland (Bishop, unpubl. data). It is unknown why larvae choose areas with high understory coverage and open canopies within a wetland, but they potentially offer greater protection from predators, higher water temperatures, higher dissolved oxygen (DO) concentrations, and possibly more prey items. All of these factors likely increase growth rates and the chances of surviving to metamorphosis.

The potential for fire exclusion to affect flatwoods salamanders negatively has been suggested (Means et al. 1996, Palis 1997b, Safer 2001). However, because no long-term monitoring datasets exist and capturing adequate numbers of individuals for research is difficult, there is limited direct evidence that suppression of wetland fires is harmful to *A. cingulatum*. Populations occasionally persist in fire-suppressed areas, but casual observations suggest that breeding areas that are fire-suppressed produce fewer larvae than those that are

We calculated the average fire return interval (i.e., the average number of years between fires) by dividing the total number of years of fire history by the number of fires. For wetlands in compartments that never burned within the observed period, we set burn frequency for the next year. For example, if a compartment containing a wetland never burned in the observed 11-year period, we set burn frequency at a default value of 12 (i.e., the area burned an average of every 12 years). Thus, mean fire return interval may slightly overestimate the actual fire frequency. However, only nine of 154 wetlands were located in compartments that never burned in the observed period. All statistical analyses were conducted in SPSS 11.0.

RESULTS

Management areas containing historical breeding wetlands (Total N = 154) had an average fire return interval (\pm SD) of 4.4 ± 2.9 years (range: 1.2–12). Fires were more than three times as common in winter and early spring (December–April) as they were

in summer (May–August) (Figure 1).

DISCUSSION

Adult flatwoods salamanders move to breeding wetlands from October–January on rainy nights associated with cold




Figure 1. Percentage of fires (Total N = 228) by month over an 11-year period in locations containing flatwoods salamander breeding wetlands.

burned frequently (D. Printiss, Apalachicola National Forest, pers. comm.). In the absence of direct evidence, we must rely on indirect evidence to determine appropriate management strategies and, whenever possible, mimic the natural processes that shaped the evolutionary history of the species.

Ephemeral depression wetlands in the southeastern coastal plain are structured by two primary factors – hydrology and fire frequency (Kirkman et al. 2000). Fire frequency declines as hydroperiod increases; however, fire frequency also may affect hydrology. Fire exclusion allows woody vegetation to increase, possibly altering evapotranspiration rates. In flatwoods communities, removal of woody vegetation can alter the hydrology of uplands and wetlands (Sun et al. 2000, Bliss and Comerford 2002). Cypress domes, limesink ponds, Carolina bays, and other southeastern ephemeral wetlands naturally burn when dry (Ewel and Mitsch 1978, Sutter and Kral 1994, Ewel 1995), but managers have been reluctant to incorporate burning wetlands in fire-management plans. As a result, many depression wetlands in the southeast likely are overgrown after years of fire suppression (Huffman and Blanchard 1991).

In other aquatic ecosystems, canopy overgrowth has led to local population extinctions (Skelly et al. 1999) and loss of diversity (Werner and Glennemeier 1999) of amphibians. Closed canopies diminish growth and developmental and survivorship rates in some species, presumably because shading decreases the amount of understory vegetation, lowers DO concentration, limits prey base, and decreases water temperature (Werner and Glennemeier 1999, Skelly et al. 2002, Halverson et al. 2003). Larval amphibians are highly sensitive, both behaviorally and physiologically, to changes in DO and water temperature in the field and lab (Costa 1967, Noland and Ultsch 1981, Nie et al. 1999). Fires also affect nutrient and pH levels in depression wetlands in longleaf pine ecosystems (Battle and Golladay 2003), factors that may alter habitat composition and developmental rates of larvae. Amphibian species that breed in ephemeral habitats must complete

metamorphosis before dry-down; hence, growth and survivorship rates need to be maximized. Slow-growing individuals often metamorphose at smaller sizes, a factor that may affect adult fitness in *Ambystoma* (Semlitsch et al. 1988, Taylor and Scott 1997).

Land managers recognize that fire is a critical component in maintaining the longleaf ecosystem, but there is considerable debate on the importance of seasonality of fires. Some suggest that growing-season fires are better than winter fires at maintaining the longleaf vegetation community (e.g., Brockway and Lewis 1997), but others believe that fire frequency is the most important characteristic, rather than season (e.g., Hiers et al. 2000). Too often, the number of acres burned, the frequency of fires, or the seasonality, rather than the ecological effects, evaluates the success of a prescribed burning program.

In the Southeast, prescribed fires typically are applied in winter and early spring, principally because winter weather conditions often are more favorable for controlling burns than summer fires (Wade and Lunsford 1989). Winter and early spring fires, however, are unlikely to burn through water-filled wetlands, except in drought years (Battle and Golladay 2003, pers. observations). Land managers in many locations make an effort to apply growing-season fires, although months that are considered part of the growing season do not necessarily correspond to the peak lightning season (May-August, Robbins and Myers 1992). Fire personnel may be reluctant to burn wetlands because of the misconception they are protecting them or their desire to avoid ‘muck’ fires, so firebreaks are plowed around the perimeters of wetlands. Water-filled wetlands often are used intentionally as natural firebreaks in prescribed burns.

Besides excluding fires, we have other concerns about firebreaks surrounding wetlands. First, they may alter hydrology. Second, adult flatwoods salamanders deposit eggs terrestrially prior to inundation (Anderson and Williamson 1976). The environmental cues that a female uses to decide where to deposit her eggs are un-

known. If females search for a depression area in the general vicinity of the wetland, a plow line may be sufficient to trigger oviposition. Lastly, firebreaks are potential death traps to larval amphibians. When a wetland is filled completely, a firebreak may be submerged within the boundaries of the wetland. As the wetland dries, the plow line becomes an isolated moat, typically drying before the interior of the wetland and leaving any trapped amphibians with less time to complete metamorphosis. In January 2002, we counted more than 500 dead leopard frog (*Rana sphenoccephala utricularia*) tadpoles in a recently dried plow line surrounding one *A. cingulatum* breeding wetland.

The accumulation of organic material in fire-suppressed wetlands concerns fire personnel wishing to avoid lingering fires and resulting smoke (Miller et al. 1998). An increase in organic material may contribute to successional change within the wetland (Russell et al. 1999) and alter DO concentration (Werner and Glennemeier 1999). This is not a concern in many *A. cingulatum* breeding wetlands because of frequent dry periods (e.g., Battle and Golladay 2001); however, some have substantial organic layers. Continuing to exclude fires, however, is not a solution and only allows organic matter to continue to accumulate, possibly leading to catastrophic fires in the future in addition to potentially modifying the habitat of larval flatwoods salamanders. Even if fires only burn through the outer edge of wetlands, larval salamanders likely will benefit from greater understory growth, higher water temperatures, and higher dissolved oxygen levels. Larvae often are captured in the inundated wiregrass ecotone (pers. observations), which flowers after growing-season fires (Seamon et al. 1989).

It is difficult to know the appropriate amount of canopy and understory coverage in ephemeral depression wetlands, if such a range exists. Likewise, we do not know what the natural fire frequency in wetland basins should be. The preferred fire frequency may vary among different amphibian species (e.g., Schurbon and Fauth 2003). If the longleaf pine ecosystem historically burned every 1-4 years

(Clewell 1989, Stout and Marion 1993, Frost 1995) and primarily in summer, fires frequently would have burned through ephemeral wetlands. The mean fire return interval for management areas surveyed in this study was 4.4 years in known breeding areas. However, fires were applied primarily in winter and early spring, and some compartments containing breeding wetlands were never burned in the 11-year period. Because breeding wetlands typically are inundated during winter and early spring, they likely are being burned at an unnaturally low frequency. In addition, most private lands typically do not burn as frequently as the lands surveyed in this study, many of which are actively managed for the federally endangered red-cockaded woodpecker (*Picoides borealis* Vieillot). In the absence of certainty of what the habitat 'should' look like, the best approach may be a mixed strategy that alters the frequency and seasonality of fires (Hiers et al. 2000).

Pilliod et al. (2003) provided a thorough review of the wide-reaching effects of fire on amphibians. More long-term research is needed to understand the relationship between fire history and the abundance, distribution, and reproductive success of *A. cingulatum*. With a declining species, however, management decisions must be made now. Management strategies should be based on what is known about the biology of the species, the hydrology of their breeding wetlands, studies of other amphibian taxa with similar life histories, and the historical importance of lightning-season fires in the longleaf ecosystem. Until data suggest otherwise, we should burn these wetlands periodically. At the very least, land managers need to stop intentionally excluding fires from them. We may, in fact, be required to do so by the Endangered Species Act (White 1989).

ACKNOWLEDGMENTS

We thank all agencies that contributed fire history information. Jim Fraser, Kevin Hiers, John Palis, David Printiss, Dirk Stevenson, and three anonymous reviewers provided helpful comments on an earlier version of this manuscript. This research

was supported by a grant from the natural resources branch (Jackson Guard) at Eglin Air Force Base administered partially through the USGS Biological Resources Division, Virginia Cooperative Fish and Wildlife Research Unit.

David Bishop is a Ph.D. student in the Department of Fisheries and Wildlife Sciences at Virginia Tech. Major interests include applying behavioral research to conservation planning and improving environmental education.

Carola Haas is an Associate Professor of wildlife ecology in the Department of Fisheries and Wildlife Sciences at Virginia Tech. Her research focuses on maintenance of native wildlife populations on lands used for commodity production or other intensive human activities.

LITERATURE CITED

- Anderson, J.D., and Williamson, G.K. 1976. Terrestrial mode of reproduction in *Ambystoma cingulatum*. *Herpetologica* 32:214-221.
- Ash, A.N. 1997. Disappearance and return of plethodontid salamanders to clearcut plots in the southern Blue Ridge Mountains. *Conservation Biology* 11:983-989.
- Battle, J., and S.W. Golladay. 2001. Hydroperiod influence on breakdown of leaf litter in cypress-gum wetlands. *American Midland Naturalist* 146:128-145.
- Battle, J., and S.W. Golladay. 2003. Prescribed fire's impact on water quality of depression wetlands in southwestern Georgia. *American Midland Naturalist* 150:15-25.
- Bliss, C.M., and N.B. Comerford. 2002. Forest harvesting influence on water table dynamics in a Florida flatwoods landscape. *Soil Science Society of America Journal* 66:1344-1349.
- Brockway, D.G., and C.E. Lewis. 1997. Long-term effects of dormant-season prescribed fire on plant community diversity, structure and productivity in a longleaf pine wiregrass ecosystem. *Forest Ecology and Management* 96:167-183.
- Clewell, A.F. 1989. Natural history of wiregrass (*Aristida stricta* Michx., Gramineae). *Natural Areas Journal* 9:223-233.
- Costa, H.H. 1967. Avoidance of anoxic water by tadpoles of *Rana temporaria*. *Hydrobiologia* 30:374-384.

- deMaynadier, P.G., and M.L. Hunter, Jr. 1995. The relationship between forest management and amphibian ecology: a review of the North American literature. *Environmental Reviews* 3:230-261.
- Ewel, K.C. 1995. Fire in cypress swamps in the southeastern United States. *Proceedings of the Tall Timbers Fire Ecology Conference* 19:111-116.
- Ewel, K.C., and W.J. Mitsch. 1978. The effects of fire on species composition in cypress dome ecosystems. *Florida Scientist* 41:25-31.
- Frost, C.C. 1995. Presettlement fire regimes in southeastern marshes, peatlands, and swamps. *Proceedings of the Tall Timbers Fire Ecology Conference* 19:39-60.
- Grialou, J.A., S.D. West, and R.N. Wilkins. 2000. The effects of forest clearcut harvesting and thinning on terrestrial salamanders. *Journal of Wildlife Management* 64:105-113.
- Halverson, M.A., D.K. Skelly, J.M. Kiesecker, L.K. Freidenburg. 2003. Forest mediated light regime linked to amphibian distribution and performance. *Oecologia* 134:360-364.
- Hiers, J.K., R. Wyatt, and R.J. Mitchell. 2000. The effects of fire regime on legume reproduction in longleaf pine savannas: is season selective? *Oecologia* 125:521-530.
- Huffman, J.M., and S.W. Blanchard. 1991. Changes in woody vegetation in Florida dry prairie and wetlands during a period of fire exclusion, and after dry-growing-season fire. Pp. 75-83 in S.C. Nodvin and T.A. Waldrop, eds., *Fire in the environment: ecological and cultural perspectives*. General Technical Report SE-69, Southeastern Forest Experiment Station, Asheville, N.C.
- Kirkman, L.K. 1995. Impacts of fire and hydrological regimes on vegetation in depression wetlands of southeastern USA. *Proceedings of the Tall Timbers Fire Ecology Conference* 19:10-20.
- Kirkman, L.K., P.C. Goebel, L. West, M.B. Drew, and B.J. Palik. 2000. Depressional wetland vegetation types: a question of plant community development. *Wetlands* 20:373-385.
- Knapp, S.M., C.A. Haas, D.N. Harpole, and R.L. Kirkpatrick. 2003. Initial effects of clearcutting and alternative silvicultural practices on terrestrial salamander abundance. *Conservation Biology* 17:752-762.
- Means, D.B. 1972. Notes on the autumn breeding biology of *Ambystoma cingulatum* (Cope) (Amphibia: Urodela: Ambystomidae). *ASB Bulletin* 19:84-85.
- Means, D.B., J.G. Palis, and M. Baggett. 1996. Effects of slash pine silviculture on a Florida

- population of flatwoods salamander. *Conservation Biology* 10:426-437.
- Miller, S.J., K.J. Ponzio, M.A. Lee, L.W. Keenan, and S.R. Miller. 1998. The use of fire in wetland preservation and restoration: are there risks? Proceedings of the Tall Timbers Fire Ecology Conference 20:127-139.
- Nie, M., J.D. Crim, and G.R. Ultsch. 1999. Dissolved oxygen, temperature, and habitat selection by bullfrog (*Rana catesbeiana*) tadpoles. *Copeia* 1999:153-162.
- Noland, R., and G.R. Ultsch. 1981. The roles of temperature and dissolved oxygen in microhabitat selection by the tadpoles of a frog (*Rana pipiens*) and a toad (*Bufo terrestris*). *Copeia* 1981:645-652.
- Palis, J.G. 1996. Flatwoods salamander (*Ambystoma cingulatum* Cope). *Natural Areas Journal* 16:49-54.
- Palis, J.G. 1997a. Breeding migration of *Ambystoma cingulatum* in Florida. *Journal of Herpetology* 31:71-78.
- Palis, J.G. 1997b. Distribution, habitat, and status of the flatwoods salamander (*Ambystoma cingulatum*) in Florida, USA. *Herpetological Natural History* 5:53-65.
- Pechmann, J.H. K., D.E. Scott, J.W. Gibbons, and R.D. Semlitsch. 1989. Influence of wetland hydroperiod on diversity and abundance of metamorphosing juvenile amphibians. *Wetlands Ecology and Management* 1:3-11.
- Petranka, J.W., M.P. Brannon, M.E. Hopey, and C.K. Smith. 1994. Effects of timber harvesting on low elevation populations of southern Appalachian salamanders. *Forest Ecology and Management* 67:135-147.
- Pilliod, D.S., R.B. Bury, E.J. Hyde, C.A. Pearl, and P.S. Corn. 2003. Fire and amphibians in North America. *Forest Ecology and Management* 178:163-181.
- Robbins, L.E., and R.L. Myers. 1992. Seasonal effects of prescribed burning in Florida: a review. Miscellaneous Publication No. 8, Tall Timbers Research Station Publication, Tallahassee, Fla.
- Russell, K.R., D.H. Van Lear, and D.C. Guynn, Jr. 1999. Prescribed fire effects on herpetofaunal: review and management implications. *Wildlife Society Bulletin* 27:374-384.
- Safer, A. 2001. Natural history and ecology of the flatwoods salamander, *Ambystoma cingulatum*, on the Atlantic coastal plain. M.S. thesis, Georgia Southern University, Statesboro.
- Schurbon, J.M., and J.E. Fauth. 2003. Effects of prescribed burning on amphibian diversity in a southeastern U.S. National Forest. *Conservation Biology* 17:1338-1349.
- Seamon, P.A., R.L. Myers, L.E. Robbins, and G.S. Seamon. 1989. Wiregrass reproduction and community restoration. *Natural Areas Journal* 9:264-265.
- Sekerak, C.M., G.W. Tanner, and J.G. Palis. 1996. Ecology of flatwoods salamander larvae in breeding ponds in Apalachicola National Forest. Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies 50:321-330.
- Semlitsch, R.D., D.E. Scott, and J.H.K. Pechmann. 1988. Time and size at metamorphosis related to adult fitness in *Ambystoma talpoideum*. *Ecology* 69:184-192.
- Skelly, D.K., L.K. Freidenburg, and J.M. Kiesecker. 2002. Forest canopy and the performance of larval amphibians. *Ecology* 83:983-992.
- Skelly, D.K., E.E. Werner, and S.A. Cortwright. 1999. Long-term distributional dynamics of a Michigan amphibian assemblage. *Ecology* 80:2326-2337.
- Stout, I.J., and W.R. Marion. 1993. Pine flatwoods and xeric pine forests of the southern (lower) coastal plain. Pp. 373-447 in W.H. Martin, S.G. Boyce and A.C. Echternacht, eds., *Biodiversity of the Southeastern United States: Lowland Terrestrial Communities*. J. Wiley, New York.
- Sun, G., H. Riekerk, and L.V. Kornhak. 2000. Ground-water-table rise after forest harvesting on cypress-pine flatwoods in Florida. *Wetlands* 20:101-112.
- Sutter, R.D., and R. Kral. 1994. The ecology, status, and conservation of two non-alluvial wetland communities in the south Atlantic and eastern Gulf coastal plain, USA. *Biological Conservation* 68:235-243.
- Taylor, B.E., and Scott, D.E. 1997. Effects of larval density dependence on population dynamics of *Ambystoma opacum*. *Herpetologica* 53:132-145.
- U.S. Fish and Wildlife Service. 1999. Endangered and threatened wildlife and plants: final rule to list the flatwoods salamander as a threatened species. *Federal Register* 64:15691-15704.
- Wade, D.D., and J.D. Lunsford. 1989. A guide for prescribed fire in southern forests. Technical Publication R8-TP11, U.S. Department of Agriculture, Forest Service, Southern Region, Atlanta, Ga.
- Ware, S., C. Frost, and P.D. Doerr. 1993. Southern mixed hardwood forest: the former longleaf pine forest. Pp. 447-493 in W.H. Martin, S.G. Boyce, and A.C. Echternacht, eds., *Biodiversity of the Southeastern United States: Lowland Terrestrial Communities*. J. Wiley, New York.
- Werner, E.E., and K.S. Glennemeier. 1999. Influence of forest canopy cover on the breeding pond distributions of several amphibian species. *Copeia* 1999:1-12.
- White, D.H. 1989. Legal implications associated with use and control of fire as a management practice. Proceedings Tall Timbers Fire Ecology Conference 17:375-384.